



Plant as an Inexhaustible medicinal source of Secondary metabolites

Rathod Zalak R., Saraf Meenu S.

Department of Microbiology and Biotechnology
University School of Sciences, Gujarat University, Ahmedabad - 380009
zdurga.03@gmail.com

Abstract

There are worries about utilizing manufactured phenolic cancer prevention agents, for example, butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) as food added substances in light of the announced negative impacts on human wellbeing. In this way, a substitution of these synthetics by cancer prevention agent extractions from different nourishments has been proposed. In excess of 8000 diverse phenolic mixes have been described; foods grown from the ground are the prime wellsprings of normal cell reinforcements. So as to concentrate, measure, and distinguish bioactive mixes from a wide assortment of foods grown from the ground, scientists utilize various procedures and strategies. This survey incorporates a short depiction of a wide scope of various measures. The cancer prevention agent, antimicrobial, and anticancer properties of phenolic normal items from leafy foods are additionally talked about.

Keywords: Antioxidants; Antimicrobial, Flavonoid; Medicinal plants; Phenol.

1. INTRODUCTION

Numerous cancer prevention agent mixes can be found in foods grown from the ground including phenolics, carotenoids, anthocyanins, and tocopherols (Jakubowski, W. et al. 1997). Around 20% of realized plants have been utilized in pharmaceutical examinations, affecting the human services framework in positive manners, for example, treating malignant growth and destructive ailments (Nacz M. et al. 2006). Plants can deliver an enormous number of differing bioactive mixes. High convergences of phytochemicals, which may secure against free extreme harm, gather in products of the soil (Suffredini, I. B. 2004). Plants containing helpful phytochemicals may enhance the necessities of the human body by going about as characteristic cell reinforcements (Boots, A.W. et al. 2008; Rathod et al. 2020). Different investigations have demonstrated that numerous plants are rich wellspring of cancer prevention agents. The therapeutic significance of plants relies on the synthetically crucial and dynamic substances that produce explicit physiological activity on the human body. Flavonoids, tannin, phenolic mixes and alkaloids are the most significant bioactive parts of plants. (Khalid, S. et al. 2018) For example, nutrients A, C, E, and phenolic mixes, for example, flavonoids, tannins, and lignins, found in plants, all go about as cell reinforcements (Suffredini, I. B. 2004). The utilization of foods grown from the ground has been connected with a few medical advantages, a consequence of therapeutic properties and high dietary benefit (Valko, M. et al. 2006). An extensive number of Tolima TDF plants, among local, naturalized and presented plants, have anthelmintic movement. The families with the most noteworthy number of plants appeared in this report are Fabacea, Amaranthaceae, Annonaceae and Euphorbiaceae. The phytochemical profile of the plants with logical anthelmintic action shows that tannins, flavonoids, and alkaloids are auxiliary, prevalent metabolites in these vegetal species. (JOLA, N. J. R. et al. 2019) Eight auxiliary metabolites were recognized from the *A. pycnantha* bloom including isosalipurposide which might be liable for their particular colouration. Nine optional metabolites were disconnected from the *J. mimosifolia* D. Wear Flower including the new phenylethanoid β -D-glucopyranose. Every detached compound was dormant against microscopic organisms tried at convergence of 32 μ g/mL. (Hendra, R. et al. 2019) Antioxidants control and decrease the oxidative harm in nourishments by postponing or repressing oxidation brought about by receptive oxygen species (ROS), eventually expanding the timeframe of realistic usability and nature of these nourishments (Ames, B.N. et al. 1993). Beta carotene, ascorbic corrosive, and numerous phenolics assume dynamic jobs in deferring maturing, lessening irritation, and forestalling certain malignant growths (Duthie, S. J. et al. 1996). Expanding the utilization of leafy foods has been suggested by numerous offices and human services frameworks all through the world (Vivekanathan, D. P. et al. 2003). The target of this paper is to give a survey of phytochemical examines that have tended to separating, estimating and



recognizing bioactive mixes of plants. This audit incorporates a review of the lipid oxidation process, subtleties of plants known to be cell reinforcement and antimicrobial sources, phenolic mixes, cell reinforcements from vegetables and organic products, malignant growth avoidance, extraction strategies for phenolic mixes, seclusion and sanitization of bioactive particles, and procedures for auxiliary characterization of bioactive atoms.

2. METHODS USED FOR BIOACTIVE COMPOUND EXTRACTION, ISOLATION, AND PURIFICATION

2.1. Extraction of Phenolic Compounds Using Solvents

Scientists have studied and analyzed the impact of different types of solvents, such as methanol, hexane, and ethyl alcohol, for the purpose of antioxidant extraction from various plants parts, such as leaves and seeds. In order to extract different phenolic compounds from plants with a high degree of accuracy, various solvents of differing polarities must be used (Wong, P. Y. Y. et al. 2006). Moreover, scientists have discovered that highly polar solvents, such as methanol, have a high effectiveness as antioxidants. Anokwuru et al. reported that acetone and N, N dimethylformamide (DMF) are highly effective at extracting antioxidants, while Koffi et al. found that methanol was more effective in at a large amount of phenolic contents from walnut fruits when compared to ethanol (Ruan, Z. P. et al. 2008; Anokwuru, C. P. et al. 2011). It has been reported that ethanolic extracts of Ivorian plants extracted higher concentrations/amount of phenolics compared to acetone, water, and methanol (Koffi, E. et al. 2010). Multiple solvents have been commonly used to extract phytochemicals, and scientists usually employed a dried powder of plants to extract bioactive compounds and eliminate the interference of water at the same time. Solvents used for the extraction of biomolecules from plants are chosen based on the polarity of the solute of interest. A solvent of similar polarity to the solute will properly dissolve the solute. Multiple solvents can be used sequentially in order to limit the amount of analogous compounds in the desired yield. The polarity, from least polar to most polar, of a few common solvents is as follows: Hexane < Chloroform<Petroleum ether< Ethyl acetate < Acetone < Methanol < Water.

2.2. Microwave-Assisted Extraction (MAE)

MAE has attracted the attention of researchers as a technique to extract bioactive compounds from a wide variety of plants and natural residues (Anokwuru, C. P. et al. 2011). Microwaves have electromagnetic radiation that occurs at frequencies between 300MHz to 300GHz, and wavelengths between 1cm and 1m. These electromagnetic waves consist of both an electrical field and a magnetic field. These are described as two perpendicular fields. The first application of microwaves was to heat up objects that can absorb a part of the electromagnetic energy and convert it into heat. Commercial microwave instruments commonly use the frequency 2450MHz, which corresponds to an energy output of 600-700Watts (Ballard, T. S. et al. 2010). The advanced techniques have become available to reduce the loss of bioactive compound without increasing the extraction time. Therefore, microwave-assisted extraction is demonstrated to be a good technique in multiple fields, especially in the medicinal plant area (Kingston, H. M. et al. 1998). Moreover, this technique reduced the losses of the biochemical compounds being extracted Microwave-assisted extraction (MAE) has been used as an alternative to conventional techniques for the extraction of antioxidants because of its ability to reduce both time and extraction solvent volume (Suzara, S. et al 2013). In fact, the main objective of using MAE is to heat the solvent and extract antioxidants from plants with a lesser amount of these solvents (Ballard, T. S. et al. 2010). Li et al. reported that conventional methods using various solvents presented less antioxidant activity and phenolic content than MAE (2012). Therefore, the finding confirmed that MAE was more effective at increasing antioxidant activity by measuring ferric reducing antioxidant power (FRAP), oxygen radical absorbance capacity (ORAC), and total phenolic content (TPC). The efficiency of the microwave extraction can be changed through some factors such as extraction temperature, solvent composition, and extraction time. The extraction temperature was usually studied more than other factors due to its ability to increase the efficiency of the microwave extraction. Tsubaki et al. reported that 170°C was the most effective temperature for extracting Phenolic compounds from Chinese tea. In addition, increasing the extraction temperature beyond this point resulted in a reduced extraction yield (2000). Christophoridou et al. used a new microwave-assisted extraction (MAE) process, which converts energy to heat, thereby cooperating with solvents in order to extract a specific compound (2005). The MAE was higher in terms of saving energy, extraction time (62 s) and extraction efficiency of bioactive compound compared to CME (2 h). Leaf presented the optimum content of total phenols (250 mg GAE/g DW) and flavonoids (13.65 mg GAE/g DW). However, the anthocyanin content was most important in pericarp extract (176.50±2.17 mg Cyd 3-glu g⁻¹ DW). The antioxidant activity was important in all parts, mainly in leaves. (Bouaoudia Madi, N. et al. 2017) Williams et al. showed many advantages of MAE, including lower solvent consumption, shorter extraction times, and higher sensitivity towards target molecules (2004). Diuzheva, A. et al determine the harpagoside and the wide phenolic pattern in Harpagophytum procumbens root and its commercial food supplements, dispersive tested with liquid-liquid microextraction (DLLME), ultrasound-assisted DLLME (UA-DLLME), and sugaring-out liquid-liquid extraction (SULLE) (2018)



2.3. Ultrasonic Assisted Extraction

Ultrasound-assisted extraction (UAE) has been used in diverse applications of food-processing technology to extract bioactive compounds from plant materials (Williams, O. J. et al. 2004). Ultrasound, with levels greater than 20 kHz, is used to disrupt plant cell walls, which helps improve the solvent's ability to penetrate the cells and obtain a higher extraction yield. UAE can use a low operating temperature through processing, maintaining a high extract quality for compounds. UAE is known to be one of the easiest extraction techniques because it uses common laboratory equipment such as an ultrasonic bath. In this technique, a smashed sample is mixed with the suitable solvent and placed into the ultrasonic bath, while temperature and extraction time are controlled (Garcia-Salas, P. et al. 2010). UAE of various organic and inorganic samples can use a wide range of solvents. Common equipment used in ultrasound-assisted extraction includes an ultrasonic bath and an ultrasonic probe system. Unfortunately, ultrasonic probe has two main negative properties mainly related to experimental repeatability and reproducibility (Corrales, M. et al 2008). Tabaraki et al. noted that green technology is necessary to protect the environment from toxic substances (2011). Therefore, extraction of phenolic compounds by ultrasound has grown during recent years due to its role in reducing the amount of solvent and energy used. Corrales et al. have shown that UAE can break down plant tissue and work properly during the production process and release of active compounds in solvents with a high efficiency (2008). Results showed an increase in antioxidant activity from 187.13 $\mu\text{molTE/g DM}$ to 308 $\mu\text{molTE/g DM}$ by using UAE as an effective method to extract antioxidants from different sources. Recently, Albu et al. studied and applied the use of ultrasound to extract phenolic compounds from rosemary (2004). Multiple criteria have been compared including ultrasonic bath extractions, ultrasonic probe system, a shaking water bath at various temperatures, and different solvents to select the most efficient method. In all situations, the operation time was dramatically decreased by applying and using the ultrasonic bath and probe systems. Similar behavior was reported by Cho et al. when extracting resveratrol from grapes (2008). In another study, Barbero et al. suggested the use of ultrasound in different industries because of its positive effects in the extraction of capsaicinoids of hot peppers (2008). Moreover, the ultrasonic method had the ability to decrease the degradation of phenolics (Luque-García, J. L. et al. 2003). Mulinacci et al. compared the extraction time of phenolic compounds from strawberries with other extraction methods such as solid-liquid, subcritical water, and microwave-assisted method (2004). The results confirmed that UAE was the most effective method.

2.4. Techniques of Isolation and Purification of Bioactive Molecules from Plants

Purification and isolation of bioactive compounds from plants is a technique that has undergone new development in recent years (Altemimi, A.W. et al. 2015; Altemimi, A. et al. 2015). This modern technique offers the ability to parallel the development and availability of many advanced bioassays on the one hand, and provided precise techniques of isolation, separation, and purification on the other. The goal when searching for bioactive compounds is to find an appropriate method that can screen the source material for bioactivity such as antioxidant, antibacterial, or cytotoxicity, combined with simplicity, specificity, and speed (Mulinacci, N. et al. 2004). In vitro methods are usually more desirable than in vivo assays because animal experiments are expensive, take more time, and are prone to ethical controversies. There are some factors that make it impossible to find final procedures or protocols to isolate and characterize certain bioactive molecules. This could be due to different parts (tissues) in a plant, many of which will produce quite different compounds, in addition to the diverse chemical structures and physicochemical properties of the bioactive phytochemicals (Sarajlija, H. et al. 2012). Both the selection and the collection of plant materials are considered primary steps to isolate and characterize a bioactive phytochemical. The next step involves a retrieval of ethno botanical information to discern possible bioactive molecules. Extracts can then be made with various solvents to isolate and purify the active compounds that are responsible for the bioactivity. Column chromatographic techniques can be used for the isolation and purification of the bioactive compounds. Developed instruments such as High Pressure Liquid Chromatography (HPLC) accelerate the process of purification of the bioactive molecule. Different varieties of spectroscopic techniques like UV-visible, Infrared (IR), Nuclear Magnetic Resonance (NMR), and mass spectroscopy can identify the purified compounds (Popova, I. E. et al. 2009).

2.5. Purification of the Bioactive Molecule

Many bioactive molecules have been isolated and purified by using paper thin-layer and column chromatographic methods. Column chromatography and thin-layer chromatography (TLC) are still mostly used due to their convenience, economy, and availability in various stationary phases. Silica, alumina, cellulose, and polyamide exhibit the most value for separating the phytochemicals. Plant materials include high amounts of complex phytochemicals, which make a good separation difficult. Therefore, increasing polarity using multiple mobile phases is useful for highly valued separations. Thin-layer chromatography has always been used to analyze the fractions of compounds by column chromatography. Silica gel column chromatography and thin layer chromatography (TLC) have been used for separation of bioactive molecules with some analytical tools (Zhang, Z. et al. 2005).



2.6. Structural Clarification of the Bioactive Molecules

Determination of the structure of certain molecules uses data from a wide range of spectroscopic techniques such as UV-visible, Infrared (IR), Nuclear Magnetic Resonance (NMR), and mass spectroscopy. The basic principle of spectroscopy is passing electromagnetic radiation through an organic molecule that absorbs some of the radiation, but not all. By measuring the amount of absorption of electromagnetic radiation, a spectrum can be produced. The spectra are specific to certain bonds in a molecule. Depending on these spectra, the structure of the organic molecule can be identified. Scientists mainly use spectra produced from either three or four regions Ultraviolet (UV), Visible, Infrared (IR), radio frequency, and electron beam for structural clarification (Popova, I. E. et al. 2009).

3. LIPID OXIDATION

Lipid oxidation can occur during the processing, shipping, and storing of many foods. Lipids (such as triglycerides, sterols, and phospholipids) readily become oxidized with exposure to an oxidative environment (Kemp, W. et al. 1991). Lipid molecules, especially those carrying polyunsaturated double bonds (i.e., linolenic acids), readily undergo oxidation within foods. Oxidatively stable oil with high melting temperature is necessary for solid fat application, and thus, highly saturated seed oil (palmitic acid and stearic acid) would be suitable for this end use (Halliwell, B. et al. 1995). Soybeans provide 56% of the world's oilseed production. However, the percentage of saturated oil is very low in seed plants (about 10%), if compared to unsaturated oil (about 90%) (Clemente, T. E. et al. 2009). Palmitic acid improves the oxidative stability of soybean oil, and can also be used to produce trans-fat-free shortening, margarine, and cosmetic products. However, this saturated short-chain fatty acid is undesirable for nutrition because its consumption results in an unfavorable lipoprotein profile in blood serum (Lakhssassi, N. et al. 2017). Stearic acid does not exhibit these cholesterolemic effects on human health (Mensink, R. P. et al. 1990). Stearic acid is less likely to be incorporated into cholesterol esters and has a neutral effect on the concentration of blood serum LDL cholesterol (Kris-Etherton, P. M. 1997). Extensive research has been performed in order to increase stearic acid content oil production in the most widely consumed legume crop in the world, soybeans. By employing induced mutagenesis, seed stearic acid content was increased by up to 7 times (Byfield, G. E. et al. 2006). Lipid oxidation in food systems can be caused by oxygen free radicals or reactive oxygen species. Free radicals are molecules with one or more unpaired electrons that work independently to cause oxidation (Lakhssassi, N. et al. 2017). Reactive oxygen species are a perfect example of oxygen free radicals. Reactive oxygen species do not solely contain free radical molecules, but also some non-free radicals that can influence lipid oxidation. Examples of non-free radical reactive oxygen species are hydrogen peroxide (H_2O_2), hydrochloric acid (HCl), ozone (O_3), and molecular oxygen (O_2) (Halliwell, B. et al. 2009). Molecular oxygen can react with linoleic acid about 1450 times faster than triplet oxygen. One of the major causes of oil rancidity is molecular oxygen. Lipid oxidation caused by the chain reaction of free radicals can be illustrated in three stages: initiation, propagation, and termination include different chemical groups such as aldehydes, ketones, alcohols, acids, or hydrocarbons. These kinds of compounds can have a negative effect on the appearance, quality, and edibility of a food product by changing the texture, color, flavor, and safety of foods, or also by producing unacceptable off odors or off tastes, even negatively affecting the nutritional value (St Angelo, A. J. et al. 1996).

4. PLANTS AS A SOURCE OF ANTIOXIDANTS

Antioxidants can be defined as bioactive compounds that inhibit or delay the oxidation of molecules (Halliwell, B. et al. 2009). Antioxidants are categorized as natural or synthetic antioxidants. Some synthetic antioxidants commonly used are: BHT, BHA, propyl gallate, and tertbutylhydroquinine. Many scientists have concerns about safety because synthetic antioxidants have recently been shown to cause health problems such as liver damage, due to their toxicity and carcinogenicity. Therefore, the development of safer antioxidants from natural sources has increased, and plants have been used as a good source of traditional medicines to treat different diseases. Many of these medicinal plants are indeed good sources of phytochemicals that possess antioxidant activities. Some typical examples of common ingredients that have been used in ethnic foods are tamarind, cardamom, lemon grass, and galangal basil. These spices or herbs have been shown to contain antioxidants (Javanmardi, J. et al. 2003). Deterioration of food due to either bacterial or fungal infection has always been a major concern, causing huge losses to food industries and societies throughout the world (Javanmardi, J. et al. 2003). Moreover, the spread of food pathogens has become a major public health concern. With an increasing awareness of the negative effects of synthetic preservatives, there has been increased demand for the use of nontoxic, natural preservatives, many of which are likely to have either antioxidant or antimicrobial activities (Negi, P. S. et al. 2005; Baharlouei, A. et al. 2010). Herbs have always been used for flavor and fragrance in the food industry, and some of them have been found to exhibit antimicrobial properties (Baharlouei, A. et al. 2011). Therefore, the call for screening and using plant materials for their antioxidant and antimicrobial properties has increased. Approximately 20% of all plant



species have been tested in both pharmacological and biological applications to confirm their safety and advantages (Suffredini, I. B. et al. 2004).

4.1. Presence of Antioxidant in Red Algae

Red algae are aquatic plant species considered one of the oldest groups of eukaryotic algae (Ruberto, G. et al. 2000). The antioxidant activity of a red alga, *Palmaria palmate*, has been studied. The results reported that 9.68 µg of ascorbic acid and 10.3 µg of total polyphenol can equally reduce activity in 1mg of dulse extracts. The reducing activity was correlated with aqueous/alcohol soluble compounds due to the presence of functional groups such as hydroxyl, carbonyl, etc., which lead to reduced or inhibited oxidation (Butterfield, N. J. et al. 2000).

4.2. Antioxidants from Monocots

Ashawat et al. studied the antioxidant properties of ethanolic extracts of *Areca catechu* and showed that *Areca catechu* had the highest antioxidant activity when compared to other eudicots like *Centella asiatica*, *Punica granatum*, and *Glycyrrhiza glabra* (2007). Londonkar and kamble studied *Pandanus odoratissimus* L. in order to determine its antioxidant activity (2009). Zahin et al. screened *Acorus calamus* to estimate antioxidant activity and total phenolic contents (2007). The observations confirmed that there was a significant correlation between the phenolic content and antioxidant activity. Another monocot, *O. sanctum*, showed that the inhibition of lipid peroxidation in vivo and in vitro increased proportionally with an increase in the concentration of the extract.

4.3. Antioxidants from Vegetables

Consumption of vegetables has been linked to a reduction in the risk of many diseases, such as cancer and cardiovascular disease, when studied in epidemiological studies (Zahin M. et al. 2007). Numerous studies have attempted to screen vegetables for antioxidant activity by using different oxidation systems. These vegetables include carrots, potatoes, sweet potatoes, red beets, cabbage, Brussels sprouts, broccoli, lettuce, spinach, onions, and tomatoes. In addition to the concise studies, which have used different methodologies to release bioactive compounds, it is becoming increasingly difficult to ignore advanced extraction methods, which have paved the way to extract bioactive compounds rapidly. Despite scientists' successes in showing the activity of vegetables' bioactive compounds, there is little known about the activity of the antioxidant components that have been isolated from these vegetables. Researchers have tended to focus on advanced methods to isolate and measure the activity of antioxidant compounds such as flavonoids, phenolic acids, tocopherols, carotenoids, and ascorbic acid (Block, G. et al. 1992).

4.4. Antioxidants from Fruits

Fruit consumption has also been linked to a reduction in the risk of many diseases (Pratt, D. E. et al. 1964). Peaches (*Prunus persica* L.) are an economically important fruit in many countries. Studies have shown that phenolic compounds found within various peach genotypes are a major source of potential antioxidants (Block, G. et al. 1992). Interestingly, peaches have shown a great inhibition of low density lipoprotein (LDL) oxidation with a percentage of antioxidant activity of 56–87%. Citrus fruits are rich sources of vitamin C (ascorbic acid), an essential nutrient with well-described antioxidant properties. However, recent studies have demonstrated that citrus also contain other bioactive compounds including flavonoids, coumarins, carotenoids, and limonoids with potential health promoting properties (Lam L. K. T. et al. 1989; Guthrie, N. et al. 2000). Accumulative evidence suggests antioxidant activities of flavonoids from a variety of plant sources (JUN YU et al. 2009). Indeed, flavonoids possess a wide range of activities in vitro (German, J. B. et al. 2000). For example, this class of bioactive compounds is known to act as free radical scavengers, to modulate enzymatic activities, and to inhibit cellular proliferation as well as possessing antibiotic, anti-allergenic, anti-diarrhea, anti-ulcer, and anti-inflammatory activities (Duthie, G. et al. 2000). Flavonoids were shown to scavenge peroxy radicals, alkyl peroxy radicals, superoxide hydroxyl radicals, and peroxy nitrite in aqueous and organic environments (Ng, T. B. et al. 2000; JUN YU et al. 2009). This antioxidant activity can be attributed to its essential compound content including hydroxycinnamic acids, chlorogenic, and neochlorogenic acids, but not to carotenoids such as *b*-carotene and *b*-cryptoxanthin. Moreover, low antioxidant activity was found in peach peel. Grape (*Vitis vinifera* L.) is a fruit crop grown throughout the world. Grapes and its juices have been studied by Plumb, G. W. et al. (1992). Phenolic compounds were high in both fresh grapes and commercial grape juices. The percentage of inhibition LDL oxidation was about 22% to 60% for fresh grapes, while it was approximately 68% to 75% for commercial grape juices, when standardized at 10mg gallic acid equivalents (GAE). According to Frankel, E. N. M. et al. (1998) both grapes and its juices exhibited high oxygen radical absorbance capacity (ORAC), and the anthocyanin pigment malvidin-3,5-diglucoside was a major compound isolated in grapes. Anthocyanins with malvidin nucleus malvidin 3-O-(6-O-p-coumaroylglucosido)-5 glucoside and phenolics were common compounds isolated from wild grapes (*Vitis coignetiae*). Wangenstein et al. tested the activity of many bioactive compounds by releasing them from grape pomace, and demonstrated that bioactive compounds have the ability to significantly inhibit LDL oxidation in the human body (2006). Grape seeds are an amazing source of polyphenol compounds including monomeric such as



catechin, epicatechin, and gallic acid, and polymeric such as procyanidins (Monagas, M. et al. 2003). Both polyphenols and carotenoids are the major phenolic compounds of apples (*Malus domestica* L.) including caffeic, quinic, and p-coumaric acids. These polyphenols can act as antioxidants. Flavanol monomers and oligomers, as well as quercetin, contribute to the beneficial health aspects of fruits and vegetables (Monagas, M. et al. 2003). Apple pomace has mainly been used as a major source of polyphenols such as chlorogenic acid (Khan, S. A. et al. 2013; Jugde, H. et al. 2008). In addition phenolics like caffeic, p-coumaroyl quinic, arbutin, p-coumaric acids, and especially flavanol procyanidins have been mentioned as constituents of apple pomace (Huang, Y. F. et al. 2012). The ability of procyanidins to work as oxygen radical scavengers, superoxides, and hydroxyl radicals was estimated. Despite the low content in total phenols in apples obtained by using acetone 70%, it has shown strong antioxidant activities towards oxidation of linoleic acid. In this case, the major bioactive compounds obtained were chlorogenic acid and phloretin glycosides; however, Vitamin C was a minor fraction in apple juice (Takos, A. M. et al. 2006). Antioxidant and antibacterial activities of various solvent (ethyl acetate, acetone, methanol, and water) extracts of *Punica granatum* peel were examined by applying the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging method. The results obtained showed a significantly higher decreasing power in the methanol extracts and a significantly higher antibacterial activity in the acetone extracts. Soong and Barlow investigated the antioxidant activity and phenolic content of various fruit seeds (Soong, Y. Y. et al. 2004). Petroleum ether was used to get rid of the excess fat from the seeds and extraction has been carried out with methanol. The 2,2-azino-bis-3 ethylbenzthiazoline-6-sulfonic acid (ABTS), DPPH, and the ferric reducing ability of plasma (FRAP) methods were used to investigate the antioxidant activity. Abdille et al. examined the antioxidant activity of *Dillenia indica* fruit using different kinds of solvents using DPPH, phospho-molybdenum, and β -carotene bleaching methods (2005). The methanol extracts showed the highest antioxidant activity, followed by the ethyl acetate and water extracts. Antioxidant activity of *Syzygium cumini* fruit in vitro has been investigated (Abdille M. H. et al. 2005). Antioxidant activity was measured by DPPH, superoxide, lipid peroxidation, and hydroxyl radical scavenging activity methods. The results brought to light a significant correlation between the concentration of the extract and the percentage of inhibition of free radicals. The antioxidant property of the fruit might be from the presence of antioxidant vitamins, anthocyanins, phenolics, and tannins. It has been reported that blackberry (*Rubus fruticosus* L.) fruit extracts produced in varying climatic regions showed that antioxidant activity depended on the genotype, rather than the climate or season (Ruan, Z. P. et al. 2008). Juntachote and Berghofer measured the stability of the antioxidant activity of ethanolic extracts for Holy basil and galangal using DPPH, superoxide, β -carotene bleaching, reducing power, and iron chelation methods (2005). They found higher antioxidant activity at neutral pH compared to an acidic pH. Holy basil and galangal extracts provided strong iron chelation activity, superoxide anion scavenging activity, and reducing power proportional to the concentration of the extracts. Liyana-Pathirana et al. investigated the antioxidant activity of cherry laurel fruit (*Laurocerasus officinalis* Roem) and its concentrated juice (Pekmez) using in vitro methods such as superoxide, DPPH scavenging activity, and inhibition of LDL oxidation (2006). The results confirmed the presence of a significantly higher antioxidant activity in pekmez compared to the cherry laurel fruit. Employing in vitro methods such as DPPH and superoxide scavenging activity, Orhan et al. measured the antioxidant activity of *Arnebia densiflora* Ledeb and observed that polar extracts had a higher antioxidant activity compared to non-polar extracts (2007). Rathee et al. studied the antioxidant activity of *Mammea longifolia* buds extracted in both methanol and aqueous ethanol. The results found a significant antioxidant activity, and the activity of aqueous ethanol was higher than methanol. The antioxidant activity of leaf extracts of *Annona* species in vitro reveals that *Annona muricata* possessed a higher antioxidant activity compared to *Annona squamosa* (Rathee, J. S. et al. 2006).

4.5. Cooking Herbs as an Important Source of Antioxidants

The antioxidant activity of 32 herbs belonging to 21 different families has been screened (Baskar, R. et al. 2007). The finding confirmed that there was a positive correlation between the total antioxidant activity and total phenolic content. Lu and Yeap Foo studied *Salvia officinalis* (L.) for its antioxidant activity and polyphenol content and reported that rosmarinic acid and various catechols were responsible for the radical scavenging activity and caffeic acid was responsible for the xanthine oxidase inhibition (Lu, Y. et al. 2001). Zhao et al. investigated the antioxidant activity of *Salvia miltiorrhiza* and *Panax notogenseng* (2006). The results showed that *Salvia miltiorrhiza* had a higher reducing power and scavenging activities against free radicals, including superoxide and hydroxyl radicals, although it showed weak hydrogen peroxide scavenging. Furthermore, Javanmardi et al. tested the Iranian *Ocimum* sp. accessions to determine the antioxidant activities and total phenolic contents and demonstrated that the antioxidant activity increased in parallel with the total phenolic content (2003). Evaluation of the pomegranate peel extracts to discover its antioxidant and antimutagenic activities using different solvents such as ethyl acetate, acetone, methanol and water has been carried out (Javanmardi J. et al. 2003). Dried extracts were examined by using the Ames test and the phosphorus molybdenum method to test both anti-mutagenic and antioxidant activities. The results showed the highest anti-mutagenic and the lowest antioxidant activity in the water extract. Moreover, the phenolic content and antioxidant activity of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) have been tested (Singh, R. P. et al. 2002). The total phenolic content was observed



to be different between parsley and cilantro leaves and stems, as well as methanol and water extracts. The methanol leaf extracts exhibited significant antioxidant activity towards both lipid and water-soluble radicals. The works also investigated the antioxidant activity of aqueous plant extracts using in vitro methods such as DPPH scavenging activity and FRAP. The results revealed a strong correlation between total antioxidant activity and phenolic content and a weak correlation between cupric ion chelators and polyphenols. The antioxidant activity and lipid peroxidation inhibition of *Satureja montana* L. subsp. *Kitaibelii* extracts were tested using hydroxyl radical scavenging. The results obtained showed that there was a significant correlation with total phenolic content (Wong, P. Y. Y. et al. 2006).

4.6. Antioxidant from Legumes

Antioxidant property of methanol extracts of *Mucuna pruriens* L. (Fabaceae) seed extracts has been investigated in vitro using the DPPH radical scavenging method. The results obtained showed a positive correlation between the antioxidant activity and the total phenolic compounds (C' etkovic', G. S. et al. 2004). Siddhuraju and Manian studied horsegram (*Macrotyloma uniflorum* Lam.) seeds to measure the antioxidant and free radical scavenging activity (2007). Acetone extracts had a higher activity of about 70% (Siddhuraju, P. et al. 2007). Samak et al. studied *Wagatea* sp. to measure its scavenging activities of superoxide and hydroxyl radicals and showed a high oxidation inhibition because it was rich in both phenolic and flavonoid contents. The authors also reported that bark and leaf extracts of *Wagatea* sp. exhibited high scavenging action against super radicals (2009).

4.7. Antioxidants from Trees

Antioxidants from trees have been also measured. Phenolics from almond hulls (*Prunus amygdalus* L.) and pine sawdust (*Pinus pinaster* L.) have been extracted employing various methods in order to determine the gram fresh yield of polyphenol compounds and antioxidant activity (Barla, A. et al. 2007). Citrus limonoids are a group of highly oxygenated triterpenoids present mainly in the Rutaceae and Meliaceae families. Research with these compounds has shown that some limonoids could induce the detoxifying enzyme glutathione S-transferase in the liver of mice and rats (Lam, L. K. T. et al. 1989). Citrus limonoids were also shown to inhibit the formation of chemically induced neoplasia in the oral cavity, forestomach, small intestine, colon, lung, and skin of laboratory animals (Lam, L. K. T. et al. 2000; Miller, E. G. et al. 2000). The limonoids can inhibit the proliferation of breast cancer cells grown in culture. However, the antioxidant activities of citrus limonoids have not yet been well-documented by JUN YU et al. (2009). An aqueous and ethanolic extracts of mango leaves were reported to be ideal antioxidants by Ling et al. (2009). The antioxidant activity was measured by the DPPH radical scavenging method. The results showed that ethanol was most appropriate either for phenolics or any bioactive compounds, while methanol was more selective for extracting polyphenolics. The antioxidant activity of juniper (*Juniperus communis*) fruit extracts has been investigated in vitro (Pinelo, M. et al. 2004). The results confirmed that both water and ethanol extract showed strong antioxidant activity. The concentration of 60µg/mL of water and ethanol extracts exhibited 84% and 92% inhibition, respectively, on the peroxidation of linoleic acid. Ibrahim et al. studied the antioxidant activity of *Cupressus sempervirens* L., and set up goals to isolate quercetin, rutin, cupress flavone, caffeic acid, and para-coumaric acid. The results showed higher antioxidant activity related to quench DPPH and identified these active compounds successively (2007). It has also been reported that Scu is more potent than kaempferol-3-O-gal in inhibiting lipid peroxidation dependent on Fe³⁺-ADP/NADPH. Neohesperidin has been reported as an alkylperoxyl radical scavenger (JUN YU et al. 2004). Higher values of antioxidant activity have been obtained by using a methanolic solvent to extract the bioactive compounds from *Anacardium occidentale*, while other solvents like ethyl acetate gave lower values of antioxidant activity (Ibrahim, N. A. et al. 2007). Kaur et al. studied the Chickrassy *Chukrasia tabularis* A. Juss leaves to confirm its ability to inhibit lipid peroxidation and showed that there was a large inhibition considering its high content of phenolic compounds (2009). *Acacia nilotica* L. antioxidant activity has been measured using ethyl acetate as a solvent to extract Phenolic compounds (Kaur, R. et al. 2009). The results exhibited the highest antioxidant activity when the concentration of extracts was relatively high.

4.8. Antioxidant from Shrubs

Many shrubs have been shown to contain antioxidant activity. Singh et al. tested several plants to measure the antioxidant activity from different extracts. The antioxidant activity was determined by using peroxide value, thiobarbituric acid, DPPH radical scavenging activity, and reducing power. The results showed that the antioxidant activity of *Coriandrum sativum* L. and *Sarcolobus globosus* L. exhibited high activity by using acetone solvent, and its activity was similar to synthetic antioxidants (Singh, G. et al. 2005). Eleven Algerian medicinal plants have been measured for phenolic compound content and antioxidant activity using the ABTS method. The tested plants showed antioxidant activity. *Artemisia campestris* L. had better antioxidant activity than caffeic acid and tocopherol. Moreover, HPLC analyses exhibited a good correlation between the antioxidant activity and hydroxycinnamic derivative content. Evaluation of *Vitex negundo* Linn seed antioxidant activity using different methods such as superoxide, hydroxyl, and DPPH scavenging activity has been carried out (Singh, G. et al. 2005).



The highest antioxidant activity was in both raw and dry heated seed extracts, while lower antioxidant activity was observed in the hydrothermally processed samples.

4.9. Characterization of Antioxidants from Other Eudicots

The nitric oxide and superoxide scavenging activity of green tea have been studied by Nakagawa and Yokozawa (Tiwari, O. P. et al. 2007), who concluded that certain tannins had the ability to exhibit excellent antioxidant activity. Zin et al. estimated the antioxidant activity of the extracts from various parts of Mengkudu (*Morinda citrifolia* L.), including the leaves, fruits, and roots, using different solvents such as methanol and ethyl acetate (2002). Ferric thiocyanate and thiobarbituric acid were used as models to observe and evaluate the antioxidant activity. The results exhibited a higher antioxidant activity in the methanol extract of Mengkudu root, although it was not significantly different from tocopherol and BHT extracts. The methanol extracts of the fruits and leaves showed unassuming activity. According to these scientists, the antioxidant activity in the roots resulted from polar and non-polar compounds, but the antioxidant activity in leaves and fruits was only due to non-polar compounds. Increase of the antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts in vitro has been shown to be proportional to the increase in the concentration of extract (Zin, Z. M. et al. 2002). Nine other extracts of Bolivian plants have been measured for radical scavenging and antioxidant activity using the DPPH and β carotene bleaching methods (Oktay, M. et al. 2003). It was found that the ethyl acetate fractions had higher radical scavenging and antioxidant activity compared to the other extracts. It has been reported that the bioactive compounds of *Rhodiola rosea* extracted in methanol showed a significant yield of phenolics, about ($153 \pm 2 \text{mg/g}$) (Parejo, I. et al. 2003). Wangensteen et al. investigated the antioxidant activity of *Ss globosus* using DPPH scavenging and inhibition of lipoxygenase (2006). Coriander had a high capacity to inhibit oxidation. There was also a positive correlation between total phenolics and antioxidant activity. Moreover, it was observed that the leaves of the coriander had higher antioxidant activity than the seeds (Parejo, I. et al. 2003). Antioxidant activity of *Phyllanthus niruri* was estimated using methanol and water as a solvent. The extracts of leaves and fruits exhibited high antioxidant activity by using the inhibition of lipid peroxidation and DPPH scavenging (Wangensteen, H. et al. 2006). The results also noticed a higher superoxide scavenging activity in the aqueous extract compared to the methanol extract. Moreover, the antioxidant and free radical scavenging activity of *Phyllanthus* species from India in an aqueous extract has been also evaluated (Harish, R. et al. 2006). The antioxidant activity was estimated using DPPH, β carotene, superoxide, nitric oxide scavenging, and reducing power methods. The extract of *Coleus aromaticus* exhibited a moderate inhibition on DPPH and nitric oxide scavenging activity. *Panax* exhibited strong iron chelating and weak superoxide scavenging. Ajila et al. carried out bioactive compounds and antioxidant activity of mango peel extract (2007). The results showed a higher concentration of anthocyanins and carotenoids in the ripe peel compared to the raw peel, while the raw peel exhibited higher polyphenol content. The range of IC₅₀ values of lipid peroxidation and DPPH were 1.39-5.24 μg of gallic acid equivalent. Chen and Yen investigated the antioxidant activity and free radical scavenging capacity of dried guava leaves and fruit (2007). The results confirmed that guava leaf and guava tea extracts had the ability to inhibit oxidation by 94-96% at a concentration of 100 $\mu\text{g/mL}$. Fruit extracts exhibited less activity compared to leaf extracts, while the scavenging effect increased with an increase in the concentration. Also, there was a correlation between antioxidant activity and phenolic compounds. Dastmalchi et al. investigated the chemical composition and antioxidant activity of water-soluble Moldavian balm (*Dracocephalum moldavica*) in vitro by using DPPH, ABTS, and superoxide activity (2007). Nitric oxide scavenging activity was calculated from *Saururus chinensis* by Marcocci et al. (1994). H_2O_2 scavenging activity was screened for various solvent of *S. chinensis* root extracts was done by Abdullah Alaklabi et al. (2018). The finding confirmed that polar compounds such as caffeic acid and rosmarinic acid were responsible for the antioxidant activity observed. Mulberry leaves were investigated to determine the antioxidant activity using different solvents (Dastmalchi, K. et al. 2007). The procedure used DPPH and inhibition of lipid peroxidation methods to evaluate its activity. The results showed that the methanolic extract exhibited the highest yield of total phenolics, and it was the most essential antioxidant in all the methods used. The antioxidant activity of kale (*Brassica oleracea* L.) has been screened after removing a fat fraction from the samples (Arabshahi-Delouee, S. et al. 2007). The extraction process used methanol to investigate its antioxidant activity while using DPPH scavenging activity as tested method. The works successfully isolated nine phenolic acids using HPLC and MS, and confirmed that the total phenolic content was correlated with DPPH scavenging activity. In another study, ethanol has been used to estimate the antioxidant activity of sun-dried cashew nuts (*Anacardium occidentale* L.) skin (Ayaz, F.A. et al. 1976). First, bioactive compounds were extracted with a protocol including lipid peroxidation, ABTS, and DPPH methods to measure the capability to inhibit oxidation. The results found that epicatechin was the major polyphenol in the extract, which was responsible for antioxidant activity. Kaviarasan et al. measured the antioxidant and antiradical activity of fenugreek (*Trigonella foenum ssp. graecum*) seeds in vitro; the results showed that there was a positive relationship between the antiradical activity and phenolic compound content in the extract (2007). Hexane and methanol were used to extract the bioactive compounds and measured the antioxidant activity of *Pueraria tuberosa* by using ABTS, lipid peroxidation, and superoxide and hydroxyl scavenging activity. An independent study has shown an inhibition of



the lipid peroxidation (Pandey, N. et al. 2010). The rhizome of the lotus (*Nelumbo nucifera* Gaertn.) has been measured for its antioxidant activity in various solvent extracts using β -Carotene bleaching and DPPH methods (Pandey, N. et al. 2010). Methanol extraction had a higher DPPH scavenging activity than acetone. *Helichrysum pedunculatum* has been tested to determine the antioxidant activity, and total phenolic and flavonoid content (Yang, D. et al. 2007). The results demonstrated that whenever the amount of phenolic content and flavonoid content was increased, higher antioxidant activity was obtained. Meot-Duros and Magn screened the leaves of *Crithmum maritimum* to show if there was any correlation between the antioxidant activity and Phenolic content and found a significant correlation between antioxidant activity and phenolic content when methanol was used as the solvent (2009). Another dicot, *Tricholepis glaberrima* L. (Asteraceae), has been investigated for antioxidant activity using different kinds of extracts (Meot-Duros, L. et al. 2009). Higher antioxidant activity was found by methanol, and a lower antioxidant activity in both chloroform and aqueous extracts. Sakat et al. investigated *Oxalis corniculata* L. in order to measure the antioxidant and anti-inflammatory activity employing methanol as a solvent. The IC₅₀ values of DPPH and nitric acid were about 93 and 73.07 $\mu\text{g/mL}$, respectively (2010). Jain et al. studied *Tabernaemontana divaricata* L. to determine the phytochemical and free radical scavenging activities in vitro. The results indicated that the antioxidant activity was the same in both ethanol and water extracts, but less in petroleum ether (2010). It has been reported that Asclepiadaceae and Periplocoideae presented high antioxidant activity, with the presence of a strong correlation between antioxidant activity and phenolic content (Jain, S. et al. 2010). Laitonjam and Kongbrailatpam studied the chemical composition and antioxidant activities of *Smilax lanceifolia* by isolating the flavonol glycoside and steroidal saponin, which showed high antioxidant activity (2010). Spinach (*Spinacea olerace* L.) is among the most popular vegetables in the world. It was domesticated and first cultivated in West Asia. According to analytical chemistry, spinach is a source of violaxanthin and neoxanthin antioxidants that cannot be commercially produced (Inbaraj, B. S. e al. 2006). Although they may be present, pigments such as carotenoids can be masked by chlorophyll in greenish vegetables such as spinach (Gandul-Rojas, B. et al. 1999). B-carotene, lutein, violaxanthin, and neoxanthin are the major carotenoids in raw spinach (Ren, D. et al. 2008). Pumpkins belong to the family Cucurbitaceae. This family is classified depending on the texture and shape of stems, such as in *Cucurbita pepo*, *Cucurbita moschata*, *Cucurbita maxima*, and *Cucurbita mixta*. Nowadays, the market offers a wide variety of vegetables, with pumpkin being one of them because of its many applications for nutrition or decoration (Alves-Rodrigues, A. et al. 2004).

5. PLANTS VITAMINS AND PHENOLIC COMPOUNDS AS ANTIOXIDANTS

5.1. Phenolic Compounds

5.1.1. Phenols and Phenolic Acid

Phenolic acids contain carboxylic acid in the chemical composition. Hydroxycinnamic and hydroxybenzoic acids are both main pillars of phenolic acids. Moreover, scientists have noted that p-coumaric, caffeic, ferulic, and sinapic acids are main components of the hydroxycinnamic acids.

5.1.2. Flavonoids

Flavonoids have a low molecular weight. Flavane is an example of a flavonoid. Flavane contains two benzene rings within its chemical composition. These two rings connect to each other through a pyrane ring. Flavones, isoflavones, flavonols, flavanones, anthocyanins, and proanthocyanidins are part of flavonoids according to the Flavonoid classification.

5.1.3. Anthocyanins

Anthocyanidins are a simple example of anthocyanins. Anthocyanidins consist of an aromatic ring that is linked to a heterocyclic ring. Moreover, the heterocyclic ring is connected to the third aromatic ring through a carbon bond (Altemimi, A. et al. 2017). Scientists have noted that anthocyanins are often found in a glycoside form. Moreover, many kinds of anthocyanins are found in nature, making these kinds of phenolic compounds very complex. Scientists have noted that anthocyanins in different kinds of fruit are considered an essential compound that can enrich and increase antioxidant activity.

5.1.4. Tannins

Tannins are natural products present in several plant families, and have large amounts of phenolic rings in the structure. Tannins are classified into two groups: hydrolyzable and condensed. Condensed tannins contain flavonoids units with several degrees of condensation. Hydrolyzable tannins contain flavonoids units with several degrees of condensation. Hydrolyzable tannins are considered a mixture of simple phenols with ester linkages in its structure. There are many factors such as alkaline compounds, mineral acids, and enzymes that have the ability to hydrolyze tannins.



5.2. Vitamins Role in Cancer Prevention

Cancer has been increasing throughout the world. It is the main cause of mortality from year to year. There were 10.4 million new cancer cases registered in 2015, and scientists predict that the number of cancer cases per year will double by 2030 (Vermerris, W. et al. 2006). Recently, many studies have shown rigorous evidence that hydroxyl radicals and the superoxide anion are involved in the development of cancer because they are biological reactive oxygen species. Compounds with high reactive oxygen species reduction activity are likely able to prevent cancer's occurrence (Hursting, S. D. et al. 1999). As shown previously, fruits and vegetables are the primary source of natural antioxidants, consisting of different kinds of antioxidant compounds such as Vitamin C, Vitamin E, carotenoids, lutein, and lycopene. Some researchers have confirmed that phenolic compounds and polyphenols are secondary plant metabolites, which are considered the best scavengers to prevent the production of free radicals. The United States has an amazing diversity of plant species. Some of them have been used for traditional medicines for a long period of time because of their various desirable activities. Kiwi and pomegranate plants extracts were screened to show the cytotoxic effects on two tumor cell lines (L20B and RD). The results have shown that the means of both L20B and RD cultures were significantly different ($p < 0.05$), and kiwi and pomegranate plant extracts exhibited a strong ability to inhibit the growth rate of L20B and RD cell lines. At concentrations of 1000 g/mL, both extracts showed a high ability to decrease the number of L20B and RD cells when compared with the control (Cai, Y. et al. 2004). The mixtures of the plant natural products have been screened in order to study their effect on human leukemia cells (Palanisamy, U. et al. 2008). The finding confirmed that mixtures of natural products were a good source for human leukemia cell inhibition. Nassr-allah et al. investigated the chemical diversity of natural products from plants in order to test their ability to work as anticancer and antioxidant agents (2009). DPPH assay was used to measure the antioxidant activity for plant extraction while using in vivo and vitro methods in order to measure the anticancer activity. The results confirmed that some natural products from Egyptian flora have the potential for use as therapeutics for diseases such as cancer (El-Shemy, H. A. et al. 2013). The effectiveness of an aqueous extract from willow leaves (*Salix safsaf*, Salicaceae) against human carcinoma cells has been tested in vivo and in vitro (Nassr-allah, A. A. et al. 2009). The findings mentioned that the metabolites for the willow extract could inhibit tumors, thereby enhancing apoptosis and causing DNA damage. The anticancer activity of different extracts from the leaves of the drumstick tree (*Moringa oleifera*) was screened in order to test against leukemia and hepatocarcinoma cells in vitro. Primary cells harvested from 10 patients with acute lymphoblastic leukemia (ALL) and 15 with acute myeloid leukemia (AML) were significantly killed by hot water and ethanol extracts. Thus, *Moringa oleifera* may have the potential for use as a natural treatment for diseases such as cancer (El-Shemy, H. A. et al. 2007). Altemimi reported that the phenolic extracts from the olive leaf extract could be used as a source of potential antioxidant and antimicrobial agents (2017).

6. PLANTS AS AN ANTIMICROBIAL SOURCE

The antibacterial activity of *Punica granatum* extracts has been investigated by using various solvents (Gil, M. I. et al. 2002). The water extract had the ability to inhibit *Bacillus subtilis* and *Staphylococcus aureus*, but the organic solvents have the ability to inhibit the growth of all the organisms tested. Shariff et al. estimated the antibacterial activity of *Rauvolfia tetraphylla* and *Physalis minima* leaves. The chloroform extract was a more powerful inhibitor of pathogenic bacteria (2006). Indian medicinal plants have been shown to have antimicrobial activity (Shariff, N. et al. 2006). About 77 extracts belonging to these plants have been tested for their antimicrobial ability against eight species of bacteria and four species of pathogenic fungi. The findings showed that water extracts of *Lantana camara* L., *Saraca asoca* L., *Acacia nilotica* L., and *Justicia zeylanica* L. caused the highest growth inhibition of all tested bacteria. The antimicrobial activity was the highest, ranging between 9.375 and 37.5g/mL and 75.0 to 300.0g/mL against both bacterial and fungal pathogens. Devi et al. investigated *Achyranthes bidentata* Blume to determine its phytochemical content and antibacterial activity (2007). The antibacterial ability of the ethanol extract effectively inhibited *Bacillus subtilis*, *Salmonella typhi*, and *Klebsiella pneumoniae*, but was less effective against *Pseudomonas* species and *Staphylococcus aureus* (Dabur, R. et al. 2007). Ethanolic extracts of *Gymnema montanum* L. have been studied to measure its antimicrobial properties against *Salmonella typhi*, *Pseudomonas aeruginosa*, and *Candida albicans* (Devi, P. U. et al. 2007). The results indicated the highest presence of antimicrobial properties in the leaf extract of *G. montanum*, correlating to its phenolic compound content. The antimicrobial activity of *Piper ribesoides* L. from methanolic root extract against *Staphylococcus aureus* has been reported (Ramkumar, K. M. et al. 2007). Interestingly, a small amount of 3.125mg/mL was enough to inhibit harmful bacteria. Leaf extracts of *Caesalpinia pulcherrima* (L.) showed higher antioxidant activity in water and ethanol extracts and lower antioxidant activity in petroleum ether extracts (Zakaria, Z. et al. 2007). *Torilis japonica* L. fruit has been observed to reduce the amount of spores, and the concentration of the vegetative cell was lower than the detection level. Ghosh et al. studied *Stevia rebaudiana* Bertoni to measure its antimicrobial properties against 10 pathogens (2008). Mahesh and Satish screened some important medicinal plants to show the antibacterial activity on human pathogenic bacteria (2008). Water and methanol were used as



solvents to extract the phenolic compounds. The finding confirmed that the methanol extract had a higher antimicrobial activity than the aqueous extract (Ghosh, S. et al. 2008). Moreover, leaf extracts of *Acacia nilotica* L., *Sida cordifolia* L., *Tinospora cordifolia* L., *Withania somnifera* L., and *Ziziphus mauritiana* L. have been studied to determine the antibacterial activity against *Bacillus subtilis*, *Escherichia coli*, *Staphylococcus aureus*, and *Pseudomonas fluorescens*, as well as studying the antifungal activity against *Aspergillus flavus*, *Dreschlera turcica*, and *Fusarium verticilloides* (Mahesh, B. et al. 2008). The highest antibacterial activity was noticed in *Acacia nilotica* and *Sida cordifolia* leaves, and the highest antifungal activity was noticed in *Acacia nilotica* bark. Water and methanol extracts of *Samanea saman* (Jacq.) exhibited a significant effect against *Xanthomonas* spp. and human pathogenic bacteria. *Pseudarthria viscida* root has been studied to measure its antimicrobial activity using ethanol as a solvent. The results showed high antimicrobial activity when compared to standard drugs like ciprofloxacin and griseofulvin. Ehsan et al. reported a high antimicrobial activity against *Staphylococcus aureus* using methanol and ethanol extracts for *Hopea parviflora* Beddome (Ehsan, B. R. et al. 2009). Ethanolic extracts of *Bryonopsis laciniosa* have been investigated for their antimicrobial activity against different Gram-positive and Gram-negative bacteria. The growth of *Staphylococcus aureus*, *Micrococcus luteus*, and *Bacillus cereus* was inhibited, as shown by a decrease in the growth zone. *Plumbago zeylanica* L. has been screened to measure the antibacterial activity in chloroform extracts to show antimicrobial activity against *Escherichia coli*, *Salmonella typhi*, and *Staphylococcus aureus* (Ehsan, B. R. et al. 2009). However, *Bacillus subtilis* and *Klebsiella* were resistant. Khond et al. studied 55 medicinal plants to measure the antimicrobial activity (2009). The higher antibacterial activities were in the extracts of *Madhuca longifolia* L., *Parkia biglandulosa* L., and *Pterospermum acerifolium* L. compared to the other plants screened. Pavithra et al. screened *Evolvulus nummularius* L. for its antibacterial activity, finding that *Escherichia coli* and *Bacillus subtilis* were the most inhibited by an ethanolic extract (Pavithra, P. S. et al. 2009). *Hygrophila spinosa* Andres leaves showed significant antibacterial activity when collected between September to October, with less activity seen during other months (Pavithra, P. S. et al. 2009). *Artemisia pallens* L. has been studied for its antimicrobial activity against seven species of bacteria (Patra, A. et al. 2009). The results found that *Bacillus cereus* was more sensitive to *A. pallens* extracts. Also, a methanolic extract exhibited higher antibacterial activity than the other solvents used. Akroum indicated the antimicrobial activity of some Algerian plants (2012). The results expressed higher antibacterial activity in methanolic extracts of *Linum capitatum*, *Camellia sinensis*, *Allium schoenoprasum*, *Vicia faba*, *Citrus paradise*, *Lippia citriodora*, *Vaccinium macrocarpon*, and *Punica granatum*. Bajpai et al. screened the antibacterial activity of *Pongamia pinnata* leaves by using methanol and ethyl acetate extracts to confirm its ability against certain pathogenic bacteria (2009). The results exhibited significant inhibition compared to streptomycin. It has been demonstrated that *Memecylon edule* has higher antibacterial activity in chloroform extracts compared to other extracts (Bajpai, V.K. et al. 2009). Gram-negative bacteria were more susceptible to the crude extracts compared to Gram-positive bacteria. Bansal et al. studied plants found in arid zones in order to determine the antibacterial efficiency (2010). An ethanolic extract of *Tinospora cordifolia* L. inhibited *Bacillus cereus* and *Staphylococcus aureus*. Kumar et al. reported *Andrographis serpyllifolia* L. to have significant antimicrobial activity against tested organisms in methanol extracts of both aerial parts and (2010). *Memecylon malabaricum*, *Cochlospermum religiosum*, and *Andrographis serpyllifolia* have been tested for their possible antimicrobial activity (Kumar, K. H. et al. 2010; Jamuna, B. A. et al. 2011). Moderate activity against both Gram-positive and Gram-negative bacteria was observed. The antimicrobial activity of an ethanolic extract of *Anethum graveolens* was better than the aqueous extract. Khanahmadi et al. (2010) found a higher antibacterial activity against Gram-positive bacteria compared to Gram-negative bacteria when an ethanolic extract of *Smyrniolum cordifolium* Boiss was used (Jamuna, B. A. et al. 2011; Khanahmadi, M. 2010). Koperuncholan et al. studied some medicinal plants of the south eastern slopes of the Western Ghats (2010). Gram-positive bacteria were more sensitive than Gram-negative bacteria to the plant extracts. Niranjana et al. screened *Schreberia swietenoides* Roxb to measure the effectiveness against human pathogenic bacteria (2010). Water and methanol extracts were most effective to prohibit growth of all the harmful bacteria tested. Different studies have isolated tannins and saponins from some Indian medicinal plants, testing the antibacterial activity against *Klebsiella pneumoniae* (Niranjana, M. H. et al. 2010; Naveen, S. M. T. 2010). Ethanol extracts of *Tinospora cordifolia* strongly inhibited *Bacillus cereus* and *Staphylococcus aureus*. Also, significant antibacterial activity from ethanolic extracts of *Coleus aromaticus* L. has been found. The most effective range of inhibition was at concentrations of 25–39 µg /mL. Vinothkumar et al. evaluated an *Andrographis paniculata* L. leaf extract's ability to inhibit the growth of Gram-positive and Gram-negative bacteria. The results found that aqueous extracts inhibited harmful microbes (2010). A positive effect of pumpkin has been observed by investigating its antimicrobial activity against *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, and *Pseudomonas aeruginosa*. Three different solvents were used to prepare the extracts: water, chloroform, and alcohol. The results showed that the alcohol extract was more powerful than both water and chloroform extracts. *Staphylococcus aureus* was sensitive to all extracts. Recently, the novel antimicrobial activity of ultrasonicated spinach leaf extracts using random amplification of polymorphic DNA (RAPD) markers and electron microscopy against both Gram-positive and Gram-negative bacteria has been revealed (Vinothkumar, P. S. K. et al. 2010). RAPD is an emerging technique used for diagnostic mutation detection within a genome. The range of the



minimum inhibitory concentrations (MICs) of the extracted leaf spinach antimicrobial substances against *Escherichia coli* and *Staphylococcus aureus* was observed between 60 and 100mg/mL. The optimal extraction conditions were at 45°C, ultrasound power of 44%, and an extraction time of 23 min. The study showed that the treated bacterial cells appeared to be damaged by a reduction in cell number. In fact, it was inferred that spinach leaf extracts exert bactericidal activity by inducing mutations in DNA and causing cell wall disruptions.

7. CONCLUSIONS

In outline, plant separates demonstrated solid cancer prevention agent limit both in vitro and in vivo, and the concentrates can be viewed as a decent wellspring of common cell reinforcements and antimicrobials. Polyphenol extraction from plants utilizing quick and proper procedures is a minimal effort technique because of the decrease in the measure of dissolvable utilized, notwithstanding dodging the requirement for longer extraction times contrasted with the regular extraction strategy. Also, characteristic bioactive mixes have been found to meddle with and forestall a wide range of malignancy. Flavonoids have been appeared to fill in as hostile to tumor (considerate, melanoma) specialists including a free radicals extinguishing instrument (i.e., OH, ROO). Truth be told, numerous examinations have demonstrated that flavonoids assume huge various jobs including mutagenic, cell harm, and cancer-causing, because of their quickening of various maturing factors. Not with standing cancer prevention agent action, the hindrance of malignant growth improvement by phenolic mixes depends on various essential cell systems. Increasingly far reaching examines identified with these mixes will upgrade pharmaceutical investigation in the field of cancer-causing illness counteraction.

CONFLICT OF INTEREST

The authors have no conflict of interest in preparing of this article.

ACKNOWLEDGEMENT

I am grateful to UGC RGNF fellowship for financial support, guide and Head of the Department of Microbiology and Biotechnology, Gujarat University for providing guidance and research facilities for conducting the research work.

REFERENCES

1. Alaklabi, A., Arif, I. A., Ahamed, A., Kumar, R. S., & Idhayadhulla, A. (2018). Evaluation of antioxidant and anticancer activities of chemical constituents of the *Saururus chinensis* root extracts. *Saudi Journal of Biological Sciences*, 25(7), 1387-1392.
2. Albu, S., Joyce, E., Paniwnyk, L., Lorimer, J. P., & Mason, T. J. (2004). Potential for the use of ultrasound in the extraction of antioxidants from *Rosmarinus officinalis* for the food and pharmaceutical industry. *Ultrasonics sonochemistry*, 11(3-4), 261-265.
3. Altemimi, A., Lakhssassi, N., Baharlouei, A., Watson, D. G., & Lightfoot, D. A. (2017). Phytochemicals: Extraction, isolation, and identification of bioactive compounds from plant extracts. *Plants*, 6(4), 42.
4. Altemimi, A., Lightfoot, D. A., Kinsel, M., & Watson, D. G. (2015). Employing response surface methodology for the optimization of ultrasound assisted extraction of lutein and β -carotene from spinach. *Molecules*, 20(4), 6611-6625.
5. Altemimi, A., Watson, D. G., Kinsel, M., & Lightfoot, D. A. (2015). Simultaneous extraction, optimization, and analysis of flavonoids and polyphenols from peach and pumpkin extracts using a TLC-densitometric method. *Chemistry Central Journal*, 9(1), 39.
6. Ame, S. N., Shigrenaga, M. K., & Hagen, T. M. (1993). Oxidants, antioxidants and degenerative diseases of ageing. *Proceedings of the national Academy of Sciences of the United States of America*, 90, 7915-7922.
7. Anokwuru, C. P., Anyasor, G. N., Ajibaye, O., Fakoya, O., & Okebugwu, P. (2011). Effect of extraction solvents on phenolic, flavonoid and antioxidant activities of three nigerian medicinal plants. *Nature and Science*, 9(7), 53-61.
8. Ashawat, M. S., Shailendra, S., & Swarnlata, S. (2007). In vitro antioxidant activity of ethanolic extracts of *Centella asiatica*, *Punica granatum*, *Glycyrrhiza glabra* and *Areca catechu*. *Res. J. Med. Plant*, 1(1), 13-16.
9. Baharlouei, A., Sharifi-Sirchi, G. R., & Bonjar, G. S. (2010). Identification of an antifungal chitinase from a potential biocontrol agent, *Streptomyces plicatus* strain 101, and its new antagonistic spectrum of activity. *Philipp. Agric. Sci*, 93, 439-445.
10. Baharlouei, A., Sharifi-Sirchi, G. R., & Bonjar, G. S. (2011). Biological control of *Sclerotinia sclerotiorum* (oilseed rape isolate) by an effective antagonist *Streptomyces*. *African Journal of Biotechnology*, 10(30), 5785-5794.



11. Ballard, T. S., Mallikarjunan, P., Zhou, K., & O'Keefe, S. (2010). Microwave-assisted extraction of phenolic antioxidant compounds from peanut skins. *Food Chemistry*, 120(4), 1185-1192.
12. Banu, K. S., & Cathrine, L. (2015). General techniques involved in phytochemical analysis. *International Journal of Advanced Research in Chemical Science*, 2(4), 25-32.
13. Barbero, G. F., Liqid, A., Palma, M., & Barroso, C. G. (2008). Ultrasound-assisted extraction of capsaicinoids from peppers. *Talanta*, 75(5), 1332-1337.
14. Beckey, H. D. (1971). *Qualitative Analyses with the Fi Mass Spectrometer*.
15. Block, G., Patterson, B., & Subar, A. (1992). Fruit, vegetables, and cancer prevention: a review of the epidemiological evidence. *Nutrition and cancer*, 18(1), 1-29.
16. Boots, A. W., Haenen, G. R., & Bast, A. (2008). Health effects of quercetin: from antioxidant to nutraceutical. *European journal of pharmacology*, 585(2-3), 325-337.
17. Bouaoudia-Madi, N., Boulekbache-Makhlouf, L., Kadri, N., Dahmoune, F., Remini, H., Dairi, S., ... & Madani, K. (2017). Phytochemical analysis of *Myrtus communis* plant: Conventional versus microwave assisted-extraction procedures. *Journal of Complementary and Integrative Medicine*, 14(4).
18. Butterfield, N. J. (2000). *Bangiomorpha pubescens* n. gen., n. sp.: implications for the evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation of eukaryotes. *Paleobiology*, 26(3), 386-404.
19. Byfield, G. E., Xue, H., & Upchurch, R. G. (2006). Two genes from soybean encoding soluble $\Delta 9$ stearoyl-ACP desaturases. *Crop science*, 46(2), 840-846.
20. Cherkaoui, A., Hibbs, J., Emonet, S., Tangomo, M., Girard, M., Francois, P., & Schrenzel, J. (2010). Comparison of two matrix-assisted laser desorption ionization-time of flight mass spectrometry methods with conventional phenotypic identification for routine identification of bacteria to the species level. *Journal of clinical microbiology*, 48(4), 1169-1175.
21. Cho, W. I., Choi, J. B., Lee, K., Chung, M. S., & Pyun, Y. R. (2008). Antimicrobial activity of torilin isolated from *Torilis japonica* fruit against *Bacillus subtilis*. *Journal of food science*, 73(2), M37-M46.
22. Christophoridou, S., Dais, P., Tseng, L. H., & Spraul, M. (2005). Separation and identification of phenolic compounds in olive oil by coupling high-performance liquid chromatography with postcolumn solid-phase extraction to nuclear magnetic resonance spectroscopy (LC-SPE-NMR). *Journal of agricultural and food chemistry*, 53(12), 4667-4679.
23. Clemente, T. E., & Cahoon, E. B. (2009). Soybean oil: genetic approaches for modification of functionality and total content. *Plant physiology*, 151(3), 1030-1040.
24. Corrales, M., Toepfl, S., Butz, P., Knorr, D., & Tauscher, B. (2008). Extraction of anthocyanins from grape by-products assisted by ultrasonics, high hydrostatic pressure or pulsed electric fields: A comparison. *Innovative Food Science & Emerging Technologies*, 9(1), 85-91.
25. Costa, S. S., Arumugamb, D., Garipeyb, Y., Rochaa, S. C., & Raghavanb, V. (2013). Spilanthal extraction using microwave: calibration curve for gas chromatography. *Chemical Engineering*, 32.
26. Diuzheva, A., Carradori, S., Andruch, V., Locatelli, M., De Luca, E., Tiecco, M., ... & Campestre, C. (2018). Use of innovative (micro) extraction techniques to characterise *Harpagophytum procumbens* root and its commercial food supplements. *Phytochemical Analysis*, 29(3), 233-241.
27. Duthie, G., & Crozier, A. (2000). Plant-derived phenolic antioxidants. *Current opinion in lipidology*, 11(1), 43-47.
28. Duthie, S. J., Ma, A., Ross, M. A., & Collins, A. R. (1996). Antioxidant supplementation decreases oxidative DNA damage in human lymphocytes. *Cancer Research*, 56(6), 1291-1295.
29. Garcia-Salas, P., Morales-Soto, A., Segura-Carretero, A., & Fernández-Gutiérrez, A. (2010). Phenolic-compound-extraction systems for fruit and vegetable samples. *Molecules*, 15(12), 8813-8826.
30. German, J. B., & Walzem, R. L. (2000). The health benefits of wine. *Annual review of nutrition*, 20(1), 561-593.
31. Guthri, N., Morley, K., Hasegawa, S., Manner, G. D., & Vandenberg, T. (2000). Inhibition of human breast cancer cells by citrus limonoids.
32. Halliwell, B., Murcia, M. A., Chirico, S., & Aruoma, O. I. (1995). Free radicals and antioxidants in food and in vivo: what they do and how they work. *Critical Reviews in Food Science & Nutrition*, 35(1-2), 7-20.
33. Hendra, R., Willis, A., & Keller, P. A. (2019). Phytochemical studies on the Australian native plant species *Acacia pycnantha* and *Jacaranda mimosifolia* D. Don. *Natural product research*, 33(14), 1997-2003.
34. Jakubowski, W., & Bartosz, G. (1997). Estimation of oxidative stress in *Saccharomyces cerevisiae* with fluorescent probes. *The international journal of biochemistry & cell biology*, 29(11), 1297-1301.
35. Javanmardi, J., Stushnoff, C., Locke, E., & Vivanco, J. M. (2003). Antioxidant activity and total phenolic content of Iranian *Ocimum* accessions. *Food chemistry*, 83(4), 547-550.
36. JOLA, N. J. R., & ESCOBAR, N. E. (2019). PROFILE OF PLANT SPECIES IN THE TROPICAL DRY FOREST OF TOLIMA (COLOMBIA) EXHIBITING ANTHELMINTIC ACTIVITY IN SHEEP. *Pak. J. Bot*, 51(5), 1737-1744.



37. Kemp, W. (1991). Infrared spectroscopy. In Organic Spectroscopy (pp. 19-99). Palgrave, London.
38. Kemp, W. (1991). Infrared spectroscopy. In Organic Spectroscopy (pp. 19-99). Palgrave, London.
39. Kemp, W. (2017). Organic spectroscopy. Macmillan International Higher Education.
40. Khalid, S., AdilShahzad, N. B., Muhammad, A., & Anwar, P. (2018). Phytochemical screening and analysis of selected medicinal plants in Gujrat. Journal of Phytochemistry and Biochemistry, 2, 108.
41. Kim, H., Moon, J. Y., Kim, H., Lee, D. S., Cho, M., Choi, H. K., ... & Cho, S. K. (2010). Antioxidant and antiproliferative activities of mango (*Mangifera indica* L.) flesh and peel. Food Chemistry, 121(2), 429-436.
42. Kingston, H. M., & Jassie, L. B. (1988). Introduction to microwave sample preparation. American Chemical Society.
43. Koffi, E., Sea, T., Dodehe, Y., & Soro, S. (2010). Effect of solvent type on extraction of polyphenols from twenty three Ivorian plants. Journal of Animal and Plant Sciences (JAPS), 5(3), 550-558.
44. Kris-Etherton, P. M., & Yu, S. (1997). Individual fatty acid effects on plasma lipids and lipoproteins: human studies. The American journal of clinical nutrition, 65(5), 1628S-1644S.
45. Kris-Etherton, P. M., Yu, S., Etherton, T. D., Morgan, R., Moriarty, K., & Shaffer, D. (1997). Fatty acids and progression of coronary artery disease. The American journal of clinical nutrition, 65(4), 1088-1090.
46. Lakhssassi, N., Colantonio, V., Flowers, N. D., Zhou, Z., Henry, J., Liu, S., & Meksem, K. (2017). Stearoyl-acyl carrier protein desaturase mutations uncover an impact of stearic acid in leaf and nodule structure. Plant physiology, 174(3), 1531-1543.
47. Lakhssassi, N., Zhou, Z., Liu, S., Colantonio, V., AbuGhazaleh, A., & Meksem, K. (2017). Characterization of the FAD2 gene family in soybean reveals the limitations of gel-based TILLING in genes with high copy number. Frontiers in plant science, 8, 324.
48. Lam, L. K., Li, Y., & Hasegawa, S. (1989). Effects of citrus limonoids on glutathione S-transferase activity in mice. Journal of Agricultural and Food Chemistry, 37(4), 878-880.
49. Li, H., Deng, Z., Wu, T., Liu, R., Loewen, S., & Tsao, R. (2012). Microwave-assisted extraction of phenolics with maximal antioxidant activities in tomatoes. Food Chemistry, 130(4), 928-936.
50. Londonkar, R., & Kamble, A. (2009). Evaluation of free radical scavenging activity of *Pandanus odoratissimus*. International journal of pharmacology, 5(6), 377-380.
51. Luque-Garcia, J. L., & De Castro, M. L. (2003). Ultrasound: a powerful tool for leaching. TrAC Trends in Analytical Chemistry, 22(1), 41-47.
52. Mensink, R. P., & Katan, M. B. (1990). Effect of dietary trans fatty acids on high-density and low-density lipoprotein cholesterol levels in healthy subjects. New England Journal of Medicine, 323(7), 439-445.
53. Mulinacci, N., Prucher, D., Peruzzi, M., Romani, A., Pinelli, P., Giaccherini, C., & Vincieri, F. F. (2004). Commercial and laboratory extracts from artichoke leaves: estimation of caffeoyl esters and flavonoidic compounds content. Journal of Pharmaceutical and Biomedical Analysis, 34(2), 349-357.
54. Naczki, M., & Shahidi, F. (2006). Phenolics in cereals, fruits and vegetables: Occurrence, extraction and analysis. Journal of pharmaceutical and biomedical analysis, 41(5), 1523-1542.
55. Negi, P. S., Chauhan, A. S., Sadia, G. A., Rohinishree, Y. S., & Ramteke, R. S. (2005). Antioxidant and antibacterial activities of various seabuckthorn (*Hippophae rhamnoides* L.) seed extracts. Food Chemistry, 92(1), 119-124.
56. Ng, T. B., Liu, F., & Wang, Z. T. (2000). Antioxidative activity of natural products from plants. Life sciences, 66(8), 709-723.
57. Popova, I. E., Hall, C., & Kubátová, A. (2009). Determination of lignans in flaxseed using liquid chromatography with time-of-flight mass spectrometry. Journal of Chromatography A, 1216(2), 217-229.
58. Rathod Z. R., Dharmal M. A., Mishra A., Patel B. V. & Saraf M. S. (2020). A Review on Phytochemical Analysis of Secondary metabolites and Extraction techniques. International Engineering Journal for Research and Development Vol.5(5), 1-10.
59. Rathod Z. R., Dharmal M. A., Sharma S., Patel B. V. & Saraf M. S. (2020). A Review on Phytochemical screening of Secondary metabolites. University Journal of Vidhya Vol.15(2), 175-190.
60. Ruan, Z. P., Zhang, L. L., & Lin, Y. M. (2008). Evaluation of the antioxidant activity of *Syzygium cumini* leaves. Molecules, 13(10), 2545-2556.
61. Ruberto, G., Baratta, M. T., Deans, S. G., & Dorman, H. D. (2000). Antioxidant and antimicrobial activity of *Foeniculum vulgare* and *Crithmum maritimum* essential oils. Planta medica, 66(08), 687-693.
62. Saračlija, H., Čukelj, N., Mršić, G. N. D., Brnčić, M., & Čurić, D. (2012). Preparation of flaxseed for lignan determination by gas chromatography-mass spectrometry method. Czech Journal of Food Sciences, 30(1), 45-52.
63. St. Angelo, A. J., Vercellotti, J., Jacks, T., & Legendre, M. (1996). Lipid oxidation in foods. Critical Reviews in Food Science & Nutrition, 36(3), 175-224.
64. Suffredini, I. B., Sader, H. S., Gonçalves, A. G., Reis, A. O., Gales, A. C., Varella, A. D., & Younes, R. N. (2004). Screening of antibacterial extracts from plants native to the Brazilian Amazon Rain Forest and Atlantic Forest. Brazilian journal of Medical and Biological research, 37(3), 379-384.



65. Tabaraki, R., & Nateghi, A. (2011). Optimization of ultrasonic-assisted extraction of natural antioxidants from rice bran using response surface methodology. *Ultrasonics sonochemistry*, 18(6), 1279-1286.
66. Tsubaki, S., Sakamoto, M., & Azuma, J. I. (2010). Microwave-assisted extraction of phenolic compounds from tea residues under autohydrolytic conditions. *Food chemistry*, 123(4), 1255-1258.
67. Urbano, M., De Castro, M. D. L., Pérez, P. M., García-Olmo, J., & Gomez-Nieto, M. A. (2006). Ultraviolet-visible spectroscopy and pattern recognition methods for differentiation and classification of wines. *Food Chemistry*, 97(1), 166-175.
68. Valko, M., Rhodes, C., Moncol, J., Izakovic, M. M., & Mazur, M. (2006). Free radicals, metals and antioxidants in oxidative stress-induced cancer. *Chemico-biological interactions*, 160(1), 1-40.
69. Vivekananthan, D. P., Penn, M. S., Sapp, S. K., Hsu, A., & Topol, E. J. (2003). Use of antioxidant vitamins for the prevention of cardiovascular disease: meta-analysis of randomised trials. *The Lancet*, 361(9374), 2017-2023.
70. Williams, O. J., RAGHAVAN, G. V., ORSAT, V., & DAI, J. (2004). Microwave-assisted extraction of capsaicinoids from capsicum fruit. *Journal of Food Biochemistry*, 28(2), 113-122.
71. Wong, P. Y., & Kitts, D. D. (2006). Studies on the dual antioxidant and antibacterial properties of parsley (*Petroselinum crispum*) and cilantro (*Coriandrum sativum*) extracts. *Food chemistry*, 97(3), 505-515.
72. Yu, J., Wang, L., Walzem, R. L., Miller, E. G., Pike, L. M., & Patil, B. S. (2005). Antioxidant activity of citrus limonoids, flavonoids, and coumarins. *Journal of agricultural and food chemistry*, 53(6), 2009-2014.
73. Zahin, M., Aqil, F., & Ahmad, I. (2009). The in vitro antioxidant activity and total phenolic content of four Indian medicinal plants. *International Journal of pharmacy and pharmaceutical Sciences*, 1(1), 88-95.
74. Zhang, Z., Pang, X., Xuwu, D., Ji, Z., & Jiang, Y. (2005). Role of peroxidase in anthocyanin degradation in litchi fruit pericarp. *Food Chemistry*, 90(1-2), 47-52.